



Araştırma Yazısı

2021; 30: 17-20

HPLC-PDA DETERMINATION OF PSEUDO TARAXASTEROL (ψ -TARAXASTEROL) IN DIFFERENT
COUSINIA SPECIES
FARKLI *COUSINIA* TÜRLERİNDE PSÖDO TARAKSASTEROL (ψ -TARAKSASTEROL) BİLEŞİĞİNİN
HPLC-PDA İLE MİKTAR TAYİNİ

Leyla PAŞAYEVA¹, Hanifa FATULLAYEV², Şehmus KILIÇ¹, Osman ÜSTÜN³, Osman TUGAY⁴

¹Erciyes University, Faculty of Pharmacy, Department of Pharmacognosy, Kayseri

²Erciyes University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, Kayseri

³Gazi University, Faculty of Pharmacy, Department of Pharmacognosy, Ankara

⁴Selçuk University, Faculty of Pharmacy, Department of Pharmaceutical Botany, Konya

ABSTRACT

Cousinia Cass. is one of the widespread genera of Asteraceae family. There is a large number of pharmacological activity studies on this genus and it has not yet been examined phytochemically in detail. Pseudo taraxasterol (ψ -taraxasterol) is a pentacyclic triterpene with different pharmacological activity such as antinociceptive, anti-inflammatory. In this report, we aimed to evaluate the ψ -taraxasterol content, which isolated from *C. stenocephala* in seven *Cousinia* species by high-performance liquid chromatography-photodiode array detector (HPLC-PDA) method. According to quantitative analyses the highest content of ψ -taraxasterol was detected in *C. davisiana*, *C. stenocephala*, and *C. ramosissima* extracts (0.423±0.004, 0.395±0.004 and 0.374±0.000 g/100g_{extract}, respectively). This is the first report on the quantification of pseudo taraxasterol by HPLC-PDA method in this genus, and this result will shed new lights on the advanced biological activity studies on species of this genus.

Keywords: Asteraceae, *Cousinia*, HPLC-PDA, pentacyclic triterpene.

ÖZ

Cousinia Cass. cinsi Asteraceae familyasına ait geniş yayılış alanına sahip cinslerden biridir. Bu cins üzerinde yok denecek kadar az farmakolojik etki ve fitokimyasal çalışmalar bulunmaktadır. Psödotaraksasterol (ψ -taraksasterol) antinosiseptif, anti-enflamatuvar aktiviteye sahip pentasikliktriterpen yapısında bileşiktir. Bu çalışmada *C. stenocephala* bitkisinden izole edilen ψ -taraksasterol bileşiğinin yedi farklı *Cousinia* türünde HPLC-PDA metodu ile miktar tayini yapılmıştır. Elde edilen sonuçlara göre enyüksek ψ -taraksasterol miktarı *C. davisiana*, *C. stenocephala* ve *C. ramosissima* ekstrel-erinde bulunmuştur (sırasıyla, 0.423±0.004, 0.395±0.004 ve 0.374±0.000 g/100g_{ekstre}). ψ -taraksasterol bileşiği bu çalışmada HPLC-PDA metodu ile bu cinste ilk defa tayin edilmiş ve bu sonuçların cinsin farklı türlerinin çeşitli biyolojik aktivite araştırmalarında faydalı olacağı düşünülmektedir.

Anahtar kelimeler: Asteraceae, *Cousinia*, HPLC-PDA, pentasiklik triterpene.

Corresponding Author: Dr.Öğr. Üyesi Leyla PAŞAYEVA, ORCID: 0000-0003-3860-7222, Department of Pharmacognosy, Faculty of Pharmacy, Erciyes University, Kayseri 38030, Turkey.

E-mail:leypasayeva@erciyes.edu.tr

Tel:+90-352-207-6666 / 28277

Öğr. Gör. Dr. Hanifa FATULLAYEV, ORCID: 0000-0002-7123-8396

Öğrenci Şehmus KILIÇ, ORCID: 0000-0002-1028-5471

Prof. Dr. Osman ÜSTÜN, ORCID: 0000-0002-6778-3834

Prof. Dr. Osman TUGAY, ORCID: 0000-0003-3980-7648

Makale Geliş Tarihi : 13.10.2019

Makale Kabul Tarihi: 09.11.2020

INTRODUCTION

Cousinia Cass. is one of the most diverse genera of Asteraceae family. This genus consists of 600-700 species distributed in Central and South-West Asia (1) and represented with six sections and 38 species in Turkey, which is 26 of them are endemic (2,3). Some species of this genera were traditionally used for the treatment of various disorders as respiratory problems, ulcers, rheumatism and inflammation (4,5). Based on previous studies cytotoxic, antioxidant, antibacterial, and hypnotic activity of different *Cousinia* species were reported (4,6-8). To date, sesquiterpene lactones (*C. Picheriana* Bornm. ex Rech.f., *C. Piptocephala* Bunge., *C. canescens* DC.), triterpenes (*C. Adenostica* Bornm.), steroids (*C. canescens* DC.), and flavonoids (*C. verbascifolia* Bunge.) have been isolated from *Cousinia* species.

The species belonging to the Asteraceae family contains various types of pentacyclic triterpenoids (e.g. oleanane, ursane, and taraxastane types). Nevertheless, due to their various pharmacological activities including anti-angiogenic, anti-inflammatory as well as antioxidant effects and the ability to enhance cell differentiation, they are widely used in medicine (9). There are some derivatives of pentacyclic triterpenes as ursane, gammacerane, lupane, and hopane (10). Among them, our main focus is on the pseudo taraxasterol. This compound is ursane type triterpene (urs-20-en-3-ol) also known as gamma-taraxasterol, heterolupeol, or calendol. The molecule formula of this compound is $C_{30}H_{50}O$ and containing six isoprene units (Figure I).




Figure I. Chemical structure of ψ -taraxasterol.

Pseudo taraxasterol mostly found in alcoholic beverages, dandelion root (*Taraxacum officinale* L.) and germinating seeds of *Calendula officinalis* L. (11). Although this compound was isolated from different plants the biological activity research on this compound is a few in literature. In a study, the anti-inflammatory and antinociceptive activity of the triterpenes β -amyrin, taraxasterol, and pseudo-taraxasterol were investigated. As a result, it was shown that the oral administration of triterpenes to mice inhibited peritoneal leukocyte infiltration. Also, the results demonstrate the anti-inflammatory and antinociceptive activity of the triterpenes were decreased when these were acetylated, while the acetylated triterpenes in mixture with myristyloxy triterpenes improved this activity (12).

In this study, we aimed to investigate qualitative and quantitative analyses of pseudo taraxasterol in seven *Cousinia* species by HPLC-PDA method.

MATERIAL AND METHOD

Chemicals

Chromatographic grade distilled water, HPLC grade methanol analytical grade formic acid was used for HPLC analyses. Pseudo taraxasterol was isolated from *C. Stenocephala* Boiss (13).

Plant Material and Preparation of Extracts

The flowering aerial parts of *C. Aintabensis* Boiss. & Hausskn., *C. davisiana* Hub.-Mor., *C. ermenekensis* Hub.-Mor., *C. foliosa* Boiss. & Bal, *C. iconica* Hub.-Mor., *C. ramosissima* DC., *C. stenocephala* Boiss. were harvested and identified by Prof. Dr. Osman Tugay. The specimens were stored at the Herbarium of Selçuk University. The details of the location and voucher number of plants were given in Table I.

Air-dried aerial parts of species (100 g) were powdered and extracted three times with methanol by maceration (during 24h), at room temperature. Combined macerates filtered and evaporated to dryness under reduced pressure at 37°C using a rotary evaporator. The crude extracts were stored in dark at -20°C. The yields of extracts were shown in Table I.

HPLC-PDA Assay

The high-performance liquid chromatographic apparatus (Shimadzu Prominence / LC-20A) with a degasser, pump (LC-20AT) and a controller coupled to a SPD-

Table I. Localities, voucher no and extract yields (%) of species.

Plant material	Localities of plants	Herbarium Voucher No	Yields (%) / Extract code
<i>C. aintabensis</i>	Mardin, Turkey; July, 2013	KNYA 11.040	10 (CA)
<i>C. davisiana</i>	Ermenek/Karaman, Konya, Turkey; July, 2013	KNYA 26.976	10 (CD)
<i>C. ermenekensis</i>	Ermenek/Karaman, Konya, Turkey; July 2013	KNYA 26.976	10 (CE)
<i>C. foliosa</i>	Ahir Dagı, Kahramanmaraş, Turkey; June, 2013	KNYA 26.977	8,7 (CF)
<i>C. iconica</i>	Ermenek/Karaman, Konya, Turkey; July 2013	KNYA 77.81	15 (CI)
<i>C. ramosissima</i>	Birecik/Şanlıurfa, Turkey; May, 2013	KNYA 26.978	10 (CR)
<i>C. stenocephala</i>	Ceylanpınar/Şanlıurfa, Turkey; July, 2013	KNYA 26.979	10 (CS)

M10Avp photodiode array detector equipped with an automatic injector interfaced to Class VP chromatography manager software. The instrument settings were performed according to the method by Sharma and Zafar (14) with slight modifications: column GL Inertsil ODS-3 (4.6mm x 250mm x 5µm), column temperature was 40°C, a flow rate of 0.8 mL/min. Detection was carried out with a sensitivity of 0.1 aufs (absorbance units full scale) between 190 and 550 nm. A mixture of methanol (A) and water: formic acid (99:1, v/v) (B) was selected as the mobile phase. The mobile phase consisted of 80% solvent A and 20% solvent B at a flow rate of 0.8 mL/min, and the injection volume was 20 µL. The retention time of the compound was 2.5 min. A 10 min equilibrium time was allowed between injections ψ -taraxasterol and sample solutions were injected three times. Compound was prepared in methanol and seven different concentration levels (0.025-1 mg/mL) were injected for the establishment of calibration curves ($y = 4642.6x - 15507$; $r^2 = 0.9984$).

RESULTS

Quantitative HPLC-PDA Analyses of Pseudo Taraxasterol

The quantitative results of the compounds are given in Table II.

Both the retention times and UV spectra were used to identify the ψ -taraxasterol. The quantities of the sub-

Table II. Content of ψ -taraxasterol in *Cousinia* extracts.

<i>Cousinia</i> species	Content*
CA	0.245±0.002
CD	0.423±0.004
CE	0.148±0.008
CF	0.214±0.005
CI	0.224±0.001
CR	0.374±0.000
CS	0.395±0.004

*Mean (g/100g_{extract}) ± SD (n=3); CA:*C. aintabensis* extract, CD:*C. davisiana* extract, CE:*C. ermenekensis* extract, CF:*C. foliosa* extract, CI:*C. iconica* extract, CR:*C. ramosissima* methanol extract, CS:*C. stenocephala* extract.

stance detected in the extracts were determined using a calibration curve. The chromatograms of ψ -taraxasterol and extracts of *Cousinia* species were shown in Figure II. As shown in the table II, the highest content of ψ -taraxasterol was found in *C. Davisiana* Hub.-Mor, *C. stenocephala* Boissand *C. ramosissima* DC extracts (0.423±0.004, 0.395±0.004 and 0.374±0.000 g/100g_{extract} respectively). This is the first report of quantitative analyses of pseudo taraxasterol in *Cousinia* species by HPLC-PDA method.

DISCUSSION

Based on chemical reports on a *Cousinia* species various chemical compounds including steroids, triterpenes, sesquiterpene lactones, and flavonoids were declared




Figure II. HPLC-PDA analyses of ψ -taraxasterol and *Cousinia* extracts with responses at 320 nm; CA: *C. aintabensis* extract, CD: *C. davisiana* extract, CE: *C. ermenekensis* extract, CF: *C. foliosa* extract, CI: *C. iconica* extract, CR: *C. ramosissima* extract, CS: *C. stenocephala* extract

(13,15-22). ψ -taraxasterol is a pentacyclic triterpene that was isolated from *Pluchea quitoc* DC., *Jatropha curcas* L., *Euphorbia nematocypha* Hand.-Mazz., *Tanacetum vulgare* L., *Achillea millefolium* L., *Kalanchoe pinnata* (Lam.) Pers., *liatris ohlingerae* (S.F.Blake) B.L.Robe, *Calotropis procera* (Aiton) Dryand., *Bryophyllum pinnatum* (Lam.) Kurz and *Sonchus arvensis* L. (23-30). Pentacyclic triterpenes have a wide spectrum of biological activities as cytotoxic, antifungal, antitumor, antiviral (31-33). To the best of our knowledge antinociceptive and anti-inflammatory effect of ψ -taraxasterol have been reported in the literature (34).

In this study, ψ -taraxasterol isolated previously from *C. stenocephala* (13) were selected and standardization of the methanol extracts of seven *Cousinia* species were achieved for the first time using a simple and efficient HPLC-PDA method. So identification and quantification of ψ -taraxasterol in *Cousinia* species will promote advanced studies that may help to protect against free radical damage and oxidative stress-related diseases. Moreover, because of the limited studies on *Cousinia* species and identified compounds, this qualitative and quantitative study combined with activity evaluation will shed new light on the advanced studies.

Acknowledgments

The authors are grateful to the Erciyes University Scientific Research Projects Coordinating Unit (BAP, project number THD-2019-8713) for financial support and Erciyes University Ziya Eren Drug Application and Research Center (ERFARMA) for providing HPLC facility.

REFERENCES

1. Reching K. *Cousinia*: morphology, taxonomy, distribution and phytogeographical implications. Proceedings of the Royal Society of Edinburgh, Section B: Biological Sciences 1986; 89, pp 45-58.
2. Davis P. Flora of Turkey and the East Aegean Islands. Edinburgh University Press, Edinburgh 1972; pp 350.
3. Güner A, Aslan S. Türkiye Bitkileri Listesi: (Damarlı Bitkiler). Nezahat Gökyiğit Botanik Bahçesi Yayınları, İstanbul 2012; s 212.
4. Amiri MS, Hosseini HA, Rajai P. Preliminary investigation on phytochemical composition and antibacterial activity of the root of *Cousinia microcarpa* Boiss. Asian J Med Pharm Res 2014; 4(4):156-159.
5. Singh KN. Traditional knowledge on ethnobotanical uses of plant biodiversity: a detailed study from the Indian western Himalaya. Biodiversity: Research and Conservation 2012; 28:63-77.
6. Sajjadi SE, Ghanadian M, Haghghi M. Isolation and identification of two phenolic compounds from a moderately cytotoxic fraction of *Cousinia verbascifolia* Bunge. Adv Biomed Res 2017; 6:66.
7. Sajjadi SE, Ghanadian M, Mouhebat L. Cytotoxic effect of *Cousinia verbascifolia* Bunge against OVCAR-3 and HT-29 cancer cells. J Herbmed Pharmacol 2015; 4(1):15-19.
8. Shahverdi AR, Khorramzadeh MR, Attar F, et al. Concomitant chemopreventive and antibacterial effects of some Iranian plants from the genus *Cousinia* (Asteraceae). Rev Bras Farmacogn 2007; 17(3):325-330.
9. Laszczyk MN. Pentacyclic triterpenes of the lupane, oleanane and ursane group as tools in cancer therapy. Planta Medica 2009; 75(15):1549-1560.
10. Patocka J. Biologically active pentacyclic triterpenes and their current medicine signification. J Appl Biomed 2003; 1(1):7-12.
11. Sethi JK, Vidal-Puig AJ. Thematic review series: adipocyte biology. Adipose tissue function and plasticity orchestrate nutritional adaptation. J Lipid Res 2007; 48(6):1253-1262.
12. da Silva FAN, de Farias Freire SM, da Rocha Borges MO, Barros FEV, da de Sousa M, de Sousa Ribeiro MN, Borges ACR. Antinociceptive and anti-inflammatory effects of triterpenes from *Pluchea quitoc* DC. aerial parts. Pharmacogn Res 2017; 9 (Suppl 1): S1.
13. Pasayeva L, Üstün O, Demirpolat E, et al. Bioactivity-guided isolation of cytotoxic and antioxidant phytochemicals from four *Cousinia* species from stenocephala bunge section. Phcog Mag 2019; 15 (65):682.
14. Sharma K, Zafar R. Simultaneous estimation of taraxerol and taraxasterol in root callus cultures of *Taraxacum officinale* Weber. Int J Pharmacogn Phytochem Res 2014; 6(3):540-546.
15. Marco JA, Sanz JF, Albiach R, Rustaiyan A, Habibi Z. Bisabolene derivatives and sesquiterpene lactones from *Cousinia* species. Phytochemistry 1993; 32 (2):395-400.
16. Plekhanova N, Turdumambetov K, Sudnitsyna I. Carbohydrates of *Cousinia*. Chem Nat Compd 1983; 19(5):603-604.
17. Rustaiyan A, Niknejad A, Sigari H, Ahmadi A. Guaianolides from *Cousinia onopordioides*. Fitoterapia 1981; 52:31-32.
18. Rustaiyan A, Sharif Z, Sadjadi AS. Two farnesol derivatives from *Cousinia adenostica*. Phytochemistry 1987; 26(9):2635-2636.
19. Turdumambetov K, Plekhanova N, Rakhimov D, Yagudaev M. Glucofructans of *Cousinia polycephala*. Chem Nat Compd 1989; 25(3):371-372.
20. Turdumambetov K, Rakhimov D, Malikova MK. Oligo- and polysaccharides from *Cousinia umbrosa*. Chem Nat Compd 2007; 43(3):308-309.
21. Ulubelen A and Tuzlacı E. Flavonoids and terpenoids from *Cousinia eriosephala*. Fitoterapia 1988; 59:350.
22. Ulubelen A, Tuzlacı E, Mericli A. Triterpenic and steroidal compounds from *Cousinia canescens*. Fitoterapia 1986; 57:269-270.
23. Chandler R, Hooper S, Hooper D, Jamieson W, Lewis E. Herbal remedies of the maritime Indians: Sterols and triterpenes of *Tanacetum vulgare* L. (Tansy). Lipids 1982; 17(2):102-106.
24. Sharma SK, Goyal N. Biological studies of the plants from genus *Pluchea*. Ann Biol Res 2011; 2 (3):25-34.
25. Xu C, Jia H-Y, Zuo B, et al. Chemical Constituents of the Aerial Parts of *Euphorbia nematocypha*. Nat Prod Commun 2016; 11(2):177-178.
26. Yadav N and Dixit V. Hepatoprotective activity of leaves of *Kalanchoe pinnata* Pers. J Ethnopharmacol 2003; 86(2-3):197-202.
27. Tiansheng L, Cantrell CL, Vargas D, et al. Terpenes from *Liatris ohlingerae*. Phytochemistry 1994; 37 (5):1295-1299.
28. Thakur S, Das P, Itoh T, Imai K, Matsumoto T. Latex extractables of *Calotropis gigantea*. Phytochemistry 1984; 23(9):2085-2087.
29. Siddiqui S, Faizi S, Siddiqui BS, Sultana N. Triterpenoids and phenanthrenes from leaves of *Bryophyllum pinnatum*. Phytochemistry 1989; 28 (9):2433-2438.
30. Hooper S, Chandler R, Lewis E, Jamieson W. Simultaneous determination of *Sonchus arvensis* L. triterpenes by gas chromatography-mass spectrometry. Lipids 1982; 17(1):60-63.
31. Chudzik M, Korzonek-Szlacheta I, Król W. Triterpenes as potentially cytotoxic compounds. Molecules 2015; 20(1):1610-1625.
32. Glinski J and Branly KL. Pentacyclic triterpenes. Patent and Trademark Office. 2001; U.S. Patent No 6,303,589.
33. Patocka J. Biologically active pentacyclic triterpenes and their current medicine signification. J Appl Biomed 2003; 1(1):7-12.
34. da Silva Fan, de Farias Freire SM, da Rocha Borges MO, et al. Antinociceptive and anti-inflammatory effects of triterpenes from *Pluchea quitoc* DC. aerial parts. Pharmacognosy Res 2017; 9(Suppl 1):1-4.